



PRESS RELEASE | PARIS | 29 SEPTEMBER 2016

The structure of the BinAB toxin revealed: one small step for Man, a major problem for mosquitoes!

Could we get rid of mosquitoes without polluting the environment? Yes, we can! The BinAB toxin, produced in crystal form by a bacterium, specifically kills the larvae of *Culex* and *Anopheles* mosquitoes, but it is inactive on tiger mosquitoes (or *Aedes*), the vectors for dengue fever and chikungunya. Knowledge of the molecular structure of BinAB is necessary if we are to broaden its spectrum of action. Having long been inaccessible, this structure is now being published on 28 September 2016 in *Nature* by an international consortium involving scientists from the Institut de Biologie Structurale (CNRS/CEA/Université Grenoble Alpes) in France, and UCLA, UCR and SLAC in the USA.

Mosquitoes are vectors for numerous devastating diseases, including malaria that is spread by *Anopheles* mosquitoes, and filariasis transmitted by *Culex* mosquitoes. The BinAB toxin, produced in the form of nanocrystals by the bacterium *Bacillus sphaericus*, specifically targets the larvae of these two groups of mosquitoes. A complex, five-step intoxication process (see insert, below) explains the environmental safety of BinAB, which is harmless to other insects, crustaceans and humans. BinAB is therefore used in many countries to regulate mosquito populations.

Unfortunately, the strength of BinAB is also its weakness: the toxin is ineffective on the larvae of *Aedes* mosquitoes, which spread the viruses for Dengue, Zika and chikungunya. A remodeling of BinAB might allow a broadening of its spectrum, but to achieve this it is necessary to understand its structure. X-ray crystallography is an excellent method to reveal the structure of a protein, but it is generally only applicable to large crystals measuring around a tenth of a millimeter. Yet, the nanocrystals of BinAB that develop *in vivo* only measure ten-thousandths of a millimeter, and once dissolved, the toxin does not recrystallize.

An international consortium of scientists led by Jacques-Philippe Colletier, CNRS scientist at the Institut de Biologie Structurale (CNRS/CEA/Université Grenoble Alpes), Brian Federici, Professor at the University of California, Riverside (UCR) and David Eisenberg, Professor at the University of California, Los Angeles (UCLA), has just published this structure, solved by working on natural nanocrystals.

Faced with the obstacle of the small size of these crystals, they employed a new type of X-ray source, a free-electron laser, delivering ultra-short but highly intense X-ray pulses. Because nothing was known of the structure of BinAB, a purely experimental approach for structure determination (*de novo* phasing) was required, which had previously only been applied to samples of known structures in order to demonstrate its feasibility.





www.cnrs.fr



Thus the structure of BinAB is not only the first to have been solved from such small crystals (~ 300 nm) but also the first unknown structure to have been revealed *de novo* using a free-electron laser. This raises hopes of solving structures from smaller and more complex natural assemblages, such as organelles, the constituents of cells.

More immediately, understanding the structure of BinAB opens the way towards broadening its spectrum of action, with the aim to develop a “three-in-one” toxin that can target the larvae of three types of mosquito: *Aedes* (in order notably to control the spread of Zika virus), *Culex* (the vector for filariasis) and *Anopheles* (the vector for malaria).



These crystals observed by electronic scanning microscopy (left) made it possible to elucidate the structure of the BinAB toxin (right). © Mari Gingery (left) / Jacques-Philippe Colletier (right).

The functioning of BinAB for the regulation of mosquito populations

The BinAB toxin is produced in the form of nanocrystals by *Bacillus sphaericus* bacteria at sporulation, or in other words when its nutrient resources diminish. Possibly attracted by the crystal, the mosquito larva eats the spore. The crystal dissolves in the larval gut where the pH is very high, releasing the BinAB toxin in a soluble form. BinAB is a binary toxin comprising two proteins, one of which specifically targets a receptor on the surface of intestinal cells (BinB), while the other serves exclusively to kill the cells (BinA). After dissolution of the crystal, BinA remains associated with BinB and the two partners are activated by the (enzymatic) digestion of their extremities (propeptides). BinB then binds to its receptor and assists the internalization of BinA – an essential step so that it can trigger the formation of a pore and thus kill the cell from the inside. What is the trophy for the bacterium? A larder where it can reproduce and survive.



www.cnrs.fr



Bibliography

De novo phasing with X-ray laser reveals mosquito larvicide BinAB structure. Jacques-Philippe Colletier, Michael R. Sawaya, Mari Gingery, Jose A. Rodriguez, Duilio Cascio, Aaron S. Brewster, Tara Michels-Clark, Robert H. Hice, Nicolas Coquelle, Sébastien Boutet, Garth J. Williams, Marc Messerschmidt, Daniel P. DePonte, Raymond G. Sierra, Hartawan Laksmono, Jason E. Koglin, Mark S. Hunter, Hyun-Woo Park, Monarin Uervirojnangkoorn, Dennis K. Bideshi, Axel T. Brunger, Brian A. Federici, Nicholas K. Sauter, David S. Eisenberg. *Nature*, 28 septembre 2016. DOI: 10.1038/nature19825 <http://www.nature.com/nature/journal/vaop/ncurrent/full/nature19825.html>

Contacts

CNRS scientist | Jacques-Philippe Colletier | T +33 (0)4 57 42 85 15 | jacques-philippe.colletier@ibs.fr
CNRS press | Véronique Etienne | T +33 (0)1 44 96 51 51 | presse@cnrs.fr